



## Control Measures for Outbreaks

### More Information

- [Prevention Measures \(/parasites/crypto/daycare/prevent.html\)](/parasites/crypto/daycare/prevent.html)
- [Control Measures for Outbreaks \(/parasites/crypto/daycare/outbreak.html\)](/parasites/crypto/daycare/outbreak.html)
- [Boil Water Advisories \(/parasites/crypto/health\\_professionals/bwa/daycare.html\)](/parasites/crypto/health_professionals/bwa/daycare.html)

### Intensified Cryptosporidiosis (Crypto) Control Measures for the Child Care Setting

Cryptosporidium is resistant to chlorine disinfection so it is tougher to kill than most disease-causing germs. The usual disinfectants, including most commonly used bleach solutions, have little effect on the parasite. An application of hydrogen peroxide seems to work best.

If an outbreak of Crypto occurs in the child care setting:

- Educate staff and parents
  - Inform all staff about the ongoing outbreak, the symptoms of Crypto, how infection is spread, and control measures to be followed.
  - Inform parents about the ongoing outbreak, the symptoms of Crypto, how infection is spread, outbreak control policies, and needed changes in hygiene and cleanliness.
  - Notify parents of children who have been in direct contact with a child or an adult caregiver with diarrhea. Parents should contact the child's healthcare provider if their child develops diarrhea.
  - Inform staff and parents of children about Crypto's potential to be a severe disease in people with weakened immune systems. Immunocompromised persons should consult their healthcare provider for further guidance.
- Exclude any child with diarrhea from the child care setting until the diarrhea has stopped.
  - Children who are infected with the parasite but who do not have diarrhea may be allowed to return.
  - Recently returning children can be grouped together in one classroom to minimize exposing uninfected children to the parasite.
  - Move adults with diarrhea to jobs that minimize opportunities for spreading infection (for example, administrative work instead of food preparation).
- Terminate all water play or swimming activities — this includes any play or activities involving water tables, [temporary inflatable or rigid fill-and-drain swimming pools and slides \(/healthywater/swimming/pools/inflatable-plastic-pools.html\)](/healthywater/swimming/pools/inflatable-plastic-pools.html), or public pool visits. The water can become contaminated and facilitate the spread of germs.
  - Exclude children diagnosed with Crypto from water-play and swimming activities for an additional 2 weeks after their diarrhea has resolved.
- Practice good hygiene.
 

Note: The hand-washing and diapering measures outlined should be routine but are especially important during outbreaks.

  - Reinforce frequent hand washing and good hand washing technique for all children and adults.
  - Good hand washing means:
    1. Wet your hands with clean running water and apply soap. Use warm water if it is available.

2. Rub hands together to make a lather and scrub all surfaces, including under the fingernails.
3. Continue rubbing hands for 20 seconds. Need a timer? Imagine singing "Happy Birthday" twice through to a friend!
4. Rinse hands well under running water.
5. Dry your hands using a disposable paper towel or a hand dryer.
6. Use your disposable paper towel, if possible, to turn off the faucet.

**Note:** Cryptosporidium is not killed by alcohol gels and hand sanitizers so these materials are of little use in controlling an outbreak.

- For children:
  - Observe hand washing or assist when needed.
  - Wash children's hands when they first arrive at the child care setting, after they use the toilet, after having their diapers changed, and before eating snacks or meals.
- For adults:
  - Wash hands after using the toilet, after helping a child use the toilet, after diapering a child, and before handling or eating food. Note: Where staffing permits, people who change diapers should not prepare or serve food.
- Reinforce good diapering practices.
  - Separate diaper-changing areas from children's play and food preparation areas.
  - Use disposable gloves and change them after each diaper change.
  - Use disposable paper over diaper changing surfaces and change it after each diaper change.
  - Ensure children wear clothing over their diapers to reduce the opportunity for leakage.
  - Wash hands: both yours and the child's after each diaper change.
- Disinfect surfaces and objects.

**Note:** The health department may instruct you to soak contaminated surfaces for 20 minutes with a 3% hydrogen peroxide (99% kill rate) and then rinse them thoroughly. No disinfectant is guaranteed to be completely effective against Cryptosporidium. However, hydrogen peroxide is more effective than standard bleach solutions.

**Note:** Do not mix hydrogen peroxide and bleach solutions. The two chemicals may react violently. In certain situations (for example, if an outbreak is caused by two or more types of germs), the health department may instruct you or a child-care facility to disinfect surfaces and objects with both hydrogen peroxide and a bleach solution. If so, disinfect with the bleach solution first and thoroughly rinse with water. Then soak with hydrogen peroxide for 20 minutes and thoroughly rinse with water.

**Note:** Hydrogen peroxide breaks down when exposed to sunlight. Store hydrogen peroxide in dedicated opaque containers — never reuse containers for a different chemical.

- Disinfect
  - Bathrooms, diaper-changing areas, and food preparation surfaces daily.
  - Toys, tabletops, and high chairs more frequently than usual (at least twice daily).
  - Dishwasher-safe toys in a commercial dishwasher that has a dry cycle or a final rinse that exceeds 113°F for 20 minutes or 122°F for 5 minutes or 162°F for 1 minute.
  - Cloth toys may be washed and heat-dried on the highest clothes dryer heat setting for

30 minutes.

- Notify the state or local health department about an excessive level of diarrhea or any Crypto cases in a daycare. Crypto is a nationally reportable disease.

## Related Links

- [Cryptosporidiosis Outbreak in a Day Care Center – Sample Letter to Day Care Center Parents !\[\]\(511a36c244659513b679df9c639945de\_img.jpg\) \(/healthywater/office/emergency/outbreak-sample-letter-child-care-parents.doc\)](/healthywater/office/emergency/outbreak-sample-letter-child-care-parents.doc) (Word - 27 KB, 2 pages)

---

Page last reviewed: May 10, 2011

Page last updated: June 16, 2011

Content source: [Centers for Disease Control and Prevention](#)

[National Center for Emerging and Zoonotic Infectious Diseases \(NCEZID\)](#)

[Division of Foodborne, Waterborne, and Environmental Diseases \(DFWED\)](#)

Notice: Linking to a non-federal site does not constitute an endorsement by HHS, CDC or any of its employees of the sponsors or the information and products presented on the site.

---

Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA  
30333, USA  
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348 - [Contact CDC-INFO](#)

