



Mechanical Wiping Increases the Efficacy of Liquid Disinfectants.

Adding wiping reduced the time to kill 6 logs of SARS-CoV-2 to 5 seconds from 5 minutes

National Microbiology Laboratory, Applied Biosafety Research Program, Safety and Environmental Services.
Public Health Agency of Canada, Winnipeg, MB



PCS 1000 Plus Microfibre Cleaning and Disinfecting

Disinfecting Clean and Disinfect Available Plastic Bucket and Lid:

Add PCS 1000 Plus Workplace Label to the pail. Add 1 liter of PCS 1000 Plus RTU or 70 mL of concentrate to 1 liter of water. Add the solution to the MT pail and place the lid loosely on top of the pail.

How to Use:

1. Remove the lid and add one PCS Hygienic Microfibre cloth to the solution. Squeeze excess liquid from the cloth so that it is just damp. It is important to maximize the absorption of pathogens and other soil. This prevents the spreading of pathogens. Place the lid loosely on top of the pail.
2. PCS 1000 Plus rapid 1-minute disinfection, combined with the PCS Hygienic Microfibre cloth's ability to trap and absorb large amounts of contamination, allows one cloth to be used versus multiple single-use disposables. The cloth maintains a high level of available chlorine even after wiping large areas. Use chlorine test strips.
3. Use the wipe-and-go process—allow to air dry.
4. One cloth can be used to oxidize, clean, and disinfect one patient space, replacing many single-use disposable wipes.
5. Do not double dip.
6. Always place the lid loosely on top of the pail.
7. PCS recommends changing solution in bucket daily. Diluted solutions in sealed containers remain effective for at least 30 days.

Manual Decontamination of Used PCS Hygienic Microfibre Cloth:

Rinse the cloth with warm water and squeeze out excess liquid. Place the cloth in PCS 1000 Plus disinfecting solution for at least ten minutes.
Rinse and dry before reusing the cloth.

1000 ppm Sodium Hypochlorite The World's Most Recommended Disinfectant for Outbreaks

